

Research in the Prediction of Solar Activity

Principal Investigator: Haimin Wang
Big Bear Solar Observatory
New Jersey Institute of Technology
Newark, NJ 07102

phone: (973) 596-5781 fax: (973) 596-5794 e-mail: haimin@solar.njit.edu

Co-Principal Investigator: Phil Goode
Big Bear Solar Observatory
40386 North Shore Lane
Big Bear City, CA 92314

phone: (909) 866-5791 fax: (909) 866-4240 e-mail: pgoode@tesla.njit.edu

Co-Principal Investigator: Hal Zirin
Solar Astronomy
264-33 Caltech
Pasadena, CA 91125

phone: (626) 395-4011 fax: (626) 395-3814 e-mail: hz@caltech.edu

Award # N000149711037
<http://bbso.njit.edu>

LONG-TERM GOAL

Our long term goal is to understand the physics and relationship among flares, filament eruptions and Coronal Mass Ejections (CMEs); to understand their impact on the magnetosphere and ionosphere of the earth; and to predict them accurately.

OBJECTIVES

We predict solar flares in our continuation of the BEARALERT program, and intend to issue more detailed alerts with general information on the state of solar activity. This will be especially important since the solar activity is approaching its maximum. Secondly, we investigate the relation between erupting filaments observed at Big Bear and Coronal Mass Ejections (CMEs) observed from Large Angle Spectroscopic Coronagraph (LASCO). If such a relationship is established, filament eruptions can be used as a proxy to predict arrival of high energy particles and geomagnetic disturbances. In addition, we have been monitoring the ultraviolet solar flux by daily issuance of our K- line index.

APPROACH

We use Big Bear magnetograms and high resolution Halpha movies to evaluate magnetic structures of active regions and issue BearAlerts when we believe that flares will occur soon; we compare Big Bear full disk Halpha movies with SOHO LASCO movies to study the relationship between filament eruptions and Coronal Mass Ejections; We obtain daily full-disk CaK images and compile our K-line index to compare with ultraviolet solar flux. People involved in the projects are: Bill Marquette (BearAlerts and Kline index), Hal Zirin (BearAlert), Phil Goode (Comparison of BBSO and LASCO

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Research in the Prediction of Solar Activity				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) New Jersey Institute of Technology,Big Bear Solar Observatory,Newark,NJ,07102				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

movies) Haimin Wang (Filament Eruption and CMEs, Temperature Structure of Erupting Filaments), Guo Yang (Filament studies), Carsten Denker (Full Disk Halpha Image Processing) and Anders Johanneson (Kline index).

WORK COMPLETED

Halpaha full disk images have been improved significantly after flat-field corrections and removal of limb darkening. They are posted on the web everyday, and become a valuable source for activity prediction by Space Environment Lab and planning of space missions like SOHO and TRACE;

- K-Line indices are on line at BBSO web page, and updated daily. Associated Kline images are archived in our ftp site;
- Filament eruption list from 1991 to 1994 has been compiled and studied; similar list from 1994 to 1998 is being compiled;
- Over 40 filament eruptions from 1996 to 1998 have been compared with LASCO movies.
- BearAlerts have been improved, and the feedback from the community is excellent.
- Polar magnetograms are obtained regularly as a supplement to full disk magnetograms from WSO and NSO, for implementing Sheeley-Wang model.

RESULTS

1. We have improved our BearAlerts, which predict solar flares and filament activation. It is more accurate and more quantitative. The success rate is above 80%. We often compare our predictions with those from NOAA. Such a success is attributed to the experience of well-trained staff and improvement of our data quality.
2. We have improved our Halpaha full disk images, which are obtained by our 2K by 2K camera, with a cadence of 1 minutes. Image are enhanced significantly (Denker et al., 1998). Such a synoptic data set is the heart of the monitoring of solar activity. Sample images can be found in <http://bbso.njit.edu>. They are being used as daily planning reference for missions like SOHO and TRACE. We have used the data to derive the solar differential rotations, statistics of mini-filament eruptions, temperature structure of prominences, and prediction of solar activity.
3. We have developed software to automatically detect filament eruptions and flares. The development is the Master thesis of one of our graduate students. Automation of the system will facilitate the prediction of the space weather, as it takes many hours to days for the magnetic cloud and high energy particles to arrive at 1AU. The major parts of the software in already in place, and will be completed by the December 1998.
4. More than 40 filament eruptions have been studied in detail, to establish the relationship between filament eruptions and CMEs. At this monen, we obtained quite confusing results: about 50% of filament eruptions are associated with CMEs, and other 50% do not. Continuing study of them may shed light on the physical reason behind it. Over 1000 filament eruptions from 1991 to 1994 have

been identified and archived at <http://solar.njit.edu>., and events after 1994 are being compiled (Wang et al., 1998).

5. We continue to obtain daily CaK images with our 1024X1024 Kodak Megaplug camera. Each day, the flat field images are obtained, so photometry data can be done with high accuracy. Over 95% of correlation between the BBSO Calcium Index and the Lyman-alpha flux has been found. Thus BBSO K index can be used as good proxy of solar irradiance.
6. To predict solar activity more accurately, Li et al. (1998) studied relationship between evolution of magnetic fields and solar flares in a super active region. They found that the occurrence of flares are closely associated with new flux emergence and excess of electric currents in one polarity.

IMPACT/APPLICATION

The prediction of solar activity is so important in many areas since magnetic eruptions can have deleterious effects on satellites, upper atmosphere communications and even the North American power grid. Thus, it is critical to have the most reliable early warning system possible. For the Navy, the important factor appears to be their effects on VLF communication and on satellites, such as those involved in communication, surveillance and the global positioning system.

TRANSITIONS

Our data on World Wide Web has been used wide in the world to monitor solar activity. Our BearAlert has helped been highly regarded in the community. The filament list and daily K-index compiled by us have been used by scientists.

RELATED PROJECTS

The PI has been awarded a Space Weather grant by NSF, which established automated early warning system of filament eruption. A computer program is being developed to automatically recognize the structure of filaments. When a filament disappears, this news will be immediately broadcast via E-mail to researchers who are interested in the project and are on our BEARALERT mailing list. The broadcast information will include the time of eruption, and the coordinates and size of the filament.

Based on the list of filament eruptions recorded by the automated program, a filament index (as a function of time) will be established

The PI also obtained a NASA SRT grant, which supports a study of comparison BBSO and Yohkoh data to understand X-ray structure associated with the filament eruptions.

PUBLICATIONS

Denker, C. Johannesson, A., Marquette, W. Goode, P.R. and Wang, H, 1998, "Synoptic Full-disk Halpha Observations at BBSO", Solar Physics, in press.

Johannesson, A. Marquette, W. and Zirin, H., 1998, "A 10-Year Set of Ca II K-Line Filtergrams", Solar Physics, 177, 265.

Li, W. Lee, C., Goode, P. , Wang, J., and Wang H., 1998, "Study of Magnetic Fields, Currents and Flares", Ap. J. submitted.

Wang, H., Komenda, A.E., Tang, F. and Zirin, H., 1998, "Filament Disappearances During the Period of September 1991 through September 1994", Solar Physics, 178, 547.

Wang, H., Chae, J., Gurman, J.B., and Kucera, T.A., 1998, "Comparison of Prominences in H α and HeII 304", Solar Physics, in press.